

Functions of long non-coding RNAs in human disease and their conservation in Drosophila development

Article (Accepted Version)

Rogoyski, Oliver M, Pueyo-Marques, Jose, Couso, Juan Pablo and Newbury, Sarah F (2017) Functions of long non-coding RNAs in human disease and their conservation in Drosophila development. *Biochemical Society Transactions*, 45 (4). pp. 895-904. ISSN 0300-5127

This version is available from Sussex Research Online: <http://sro.sussex.ac.uk/id/eprint/68277/>

This document is made available in accordance with publisher policies and may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the URL above for details on accessing the published version.

Copyright and reuse:

Sussex Research Online is a digital repository of the research output of the University.

Copyright and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable, the material made available in SRO has been checked for eligibility before being made available.

Copies of full text items generally can be reproduced, displayed or performed and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.

Functions of long non-coding RNAs in human disease and their conservation in *Drosophila* development.

Oliver M. Rogoyski¹, Jose Ignacio Pueyo¹, Juan Pablo Couso^{1,2} and Sarah F. Newbury^{1*}

¹ Brighton and Sussex Medical School, Medical Research building, University of Sussex, Falmer, Brighton, BN1 9PS, U.K.

² Centro Andaluz de Biología de Desarrollo, CSIC-Universidad Pablo de Olavide, Ctra. de Utrera, Km.1, Sevilla, 41013, Spain

*Corresponding author: s.newbury@bsms.ac.uk +44 (0)1273 877874

Abstract

Genomic analysis has found that the transcriptome in both humans and *Drosophila melanogaster* features large numbers of long non-coding RNA transcripts (lncRNAs). This recently discovered class of RNAs regulates gene expression in diverse ways, and has been involved in a large variety of important biological functions. Importantly, an increasing number of lncRNAs have also been associated with a range of human diseases, including cancer. Comparative analyses of their functions among these organisms suggest that some of their modes of action appear to be conserved. This highlights the importance of model organisms such as *Drosophila*, which shares many gene regulatory networks with humans, in understanding lncRNA function and its possible impact in human health. This review discusses some known functions and mechanisms of action of lncRNAs and their implication in human diseases, together with their functional conservation and relevance in *Drosophila* development.

Introduction

The central dogma of molecular biology as proposed by Crick in 1958, often paraphrased as “DNA encodes RNA, RNA encodes protein”, implicates RNA as a molecular intermediate in the process of protein synthesis from the relevant encoding gene. As early as the 1950s however, other roles for non-coding RNAs, such as transfer RNAs and ribosomal RNAs, have been known to be vital to biological function. This showed the central dogma to be an over-simplified, if eloquent, summary of the flow of genetic information. Since then, many other types of non-coding RNA have been shown to exist, and furthermore, to be biologically relevant. In the 1990s, several studies began investigating the biological purpose of longer non protein-coding RNAs, such as *Xist* [1], which did not fit well into the RNA classifications existing at the time. With further advances in molecular techniques suggesting that only 2% of the human genome is comprised of protein-coding genes [2], and rapidly revealing lncRNAs with biological functions (including in human diseases), the topic has become an extremely promising and popular avenue of investigation.

In this review, we have used the definition of lncRNAs as being RNA transcripts longer than 200 nucleotides, which lack a significant open reading frame (greater than 100 amino acids in length) [3]. This definition is routinely used in the annotation of the *Drosophila* and other genomes. lncRNAs are highly abundant, and are found in many organisms across different taxa, including humans, mice, *Xenopus tropicalis*, *Drosophila melanogaster*, *Schizosaccharomyces pombe*, *Saccharomyces cerevisiae*, *Caenorhabditis elegans*, *Arabidopsis thaliana*, *Medicago truncatula*, and *Zea mays* [4]. lncRNAs have been shown to regulate gene expression transcriptionally [5-8] and post-transcriptionally [9-13], and have a wide

range of cellular and molecular functions. Despite these proven non-coding functions, there exist a handful of lncRNAs that have been shown to encode small open reading frame (smORF) peptides with proven cellular functions [14-19]. Recent work has shown that lncRNAs can simultaneously display biological function as both a coding, and a non-coding RNA, for example where primary transcripts of microRNAs encode regulatory peptides [20, 21]. Additionally, ribosome profiling and bioinformatics analyses have identified the existence of thousands of lncRNAs containing putatively functional translated smORFs [19, 22-25], the extent of which may depend on developmental or tissue specific context. We have therefore used the accepted definition above, which coincides with genome annotations.

Drosophila melanogaster, the common fruit fly, is a well-established model organism for geneticists, and one in which lncRNAs are known to be abundant. With an estimated 75% of human disease-linked genes having a functional orthologue in *Drosophila*, and many basic molecular and biological functions conserved between species [26, 27], *Drosophila* are an appealing whole animal model for understanding human disease. In addition to their genetic similarities, the fly genome has been incredibly well studied and fully sequenced, with a wide range of genetic tools and gene-specific knockdown and mutant lines readily available. Combined with their low maintenance cost, short generation time, high fecundity, and compound factors lending themselves to ease of establishing genetic crosses, it is easy to see why *Drosophila* have emerged as one of the foremost systems for studying the genetic components of human disease, and have already been successfully used to dissect the roles and mechanisms of certain lncRNAs [28].

As well as the general excellence of *Drosophila* as a model organism, they stand out as particularly apt for the study of lncRNA. lncRNAs evolve rapidly, and can act as flexible scaffolds tethering together one or more functional elements [29]. *Drosophila* lncRNAs also appear to accumulate relatively few deleterious changes, due to genetic drift, compared to mammalian lncRNAs [30], and therefore can be useful in developing strategies to identify lncRNA orthologues, as shown for roX lncRNA orthologues in Drosophilid species [31]. Additionally, *Drosophila* is an excellent model system to functionally characterise lncRNA-protein complexes, for example by using the GAL4-UAS system to express lncRNAs in specific tissues or by characterising the localisation of RNA-proteins within cells (e.g. 7SK snRNA [32]).

Molecular functions and mechanisms of lncRNAs, such as their binding to protein complexes, definitively need to be tested *in vivo* in order to be well characterized. For example, *in vivo* experiments have shown that only the lncRNA transcribed in the reverse direction from the Polycomb/Trithorax response elements can bind the the Polycomb Repressive Complex 2 component Enhancer of Zeste, which provides the critical Histone Methyl Transferase activity required for transcriptional silencing. This level of understanding of such complex mechanisms and interactions would be extremely difficult to achieve without the use of a tractable *in vivo* system such as that provided by *Drosophila*.

In this review, we will be examining the emerging roles and relevance of lncRNAs using recent work illustrating their biological and molecular functions in *Drosophila*. We aim to examine these recent advances in our understanding of lncRNAs through the lens of their potential relevance to humans, and particularly human disease. By doing so, we hope to provide a concise synopsis of the topic, and demonstrate the value of using *Drosophila* as a model organism for understanding the roles of lncRNAs at molecular and cellular levels, and their implications in human disease.

Abundance and localisation of lncRNAs in the human and *Drosophila* genomes

95 According to the Ensembl database, lncRNAs comprise 7841 of the 63898 annotated genes in the human
96 genome, and 2366 of the 17559 in the *Drosophila* genome. In both species, they account for a similar and
97 substantial proportion of the entire genome (12.4% and 13.5% respectively). Although only a fraction of
98 these have been investigated experimentally, information on their sequences and loci are readily available
99 through various genomic databases, both non-specific (such as Ensembl), and dedicated non-coding RNA
100 databases (such as LNCipedia, lncRNome, and lncRNadb). Additionally, significant bioinformatic work has
101 been carried out on them in terms of their expression and conservation within and across species [33].
102 With so much information on lncRNA now available, exploring this class of genes with a thorough
103 experimental approach has become more feasible in recent years.

104
105 lncRNAs vary significantly in their distribution throughout cellular compartments, with the majority of
106 transcripts residing predominantly in the nucleus, others in the cytoplasm, and some distributed more
107 evenly between the two [34, 35]. For example, the *roX* transcripts in *Drosophila* are found in the nucleus,
108 while *yar* is cytoplasmic [35]. The localisation of lncRNAs can give clues about their function; in the case of
109 a chromatin restructuring lncRNA such as *roX1* or *roX2* it must be nuclear in order to access the chromatin.
110 Localisation of particular lncRNAs can also affect their susceptibility to suppression by RNA interference and
111 antisense oligonucleotides. An example of this is the suppression of nuclear lncRNAs *MALAT1* and *NEAT1*
112 which in humans is more efficient using antisense methods, whereas cytoplasmic lncRNAs *DANCR* and
113 *OIP5-AS1* are better suppressed with RNAi methods [35].

114
115 However, the sub-cellular localisation of the majority of lncRNAs has not been well characterised, with the
116 localisation of relatively few being experimentally visualised. Single molecule RNA fluorescence *in situ*
117 hybridisation has now been used to give high resolution data for the distribution of lncRNAs in human cells
118 [34], and a systematic investigation of lncRNA localisation has been suggested as an important next step in
119 expanding our understanding of their function; as well as a useful way to shed light on the potential
120 relevance of lncRNAs to a particular mechanism.

121 122 **lncRNA in human disease**

123
124 lncRNAs have now been implicated as important factors linked to a range of human diseases. The broad
125 range of biological functions of lncRNAs is reflected in the variety of different pathologies in which their
126 aberrant expression is thought to be a contributing factor. Many lncRNAs have been shown to either be
127 expressed at aberrant levels in cancerous cells [36-67], or their levels shown to affect the growth and
128 behaviour of cancerous cells [46, 47, 49, 50, 52-56] (Table 1). This has prompted speculation that if better
129 characterised, this class of genes may present many promising biomarkers, and even novel potential
130 therapeutic targets. We cannot comprehensively cover this topic within the scope of this review, and point
131 the reader to a comprehensive review of the topic for more information [57], but instead demonstrate this
132 point with two well documented examples, below.

133
134 *MALAT1*, a highly conserved mammalian lncRNA, has been found to be overexpressed in human
135 osteosarcoma cells and cell lines [46, 47]. It is hypothesised to function as a molecular scaffold for
136 ribonucleoprotein complexes, acting as a transcriptional regulator for certain genes. Higher levels of
137 *MALAT1* have been shown to be associated with “aggressive” cancer traits such as increased migration,
138 metastasis, and clonogenic growth in non-small cell lung cancer [36-38] pancreatic [58], and prostate
139 cancer cells [39]. Indeed, inducing a knockdown of *MALAT1* in osteosarcoma cell lines inhibited cell
140 proliferation and invasion [46, 47].

141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186

The *HOTAIR* lncRNA, transcribed from an antisense Hox gene, plays an important role in the epigenetic regulation of genes thought to be due to its interactions with the Polycomb Repressive Complex 2 (PRC2) [43, 59], although recent work has indicated that PRC2 recruitment may be a downstream consequence of gene silencing, rather than initiating it [68]. *HOTAIR* is thought to act as a molecular scaffold, and is required for histone modification of particular genes across different chromosomes. Higher levels of *HOTAIR* have been found in colorectal cancer tissues, and are associated with increased tumour invasion, metastasis, vascular invasion, advanced tumour stage, and a worse prognosis in patients [43, 44]. *HOTAIR* has since been suggested for use as a biomarker for the progression and prognosis of certain cancers [44]. A *Drosophila* homologue for *HOTAIR* has not been identified, but given the similarities in polycomb regulation between species, it is likely that a targeted search might reveal such an equivalent.

Aside from cancer, strong evidence now exists linking certain lncRNAs to certain neurological pathologies [60]. lncRNAs have been shown to be relevant factors in amyotrophic lateral sclerosis, multiple sclerosis [61, 62], Alzheimer's disease [10, 63], Huntington's disease [64, 65], and Parkinson's disease, among others. For example, the *BACE1* antisense transcript (*BACE1-AS*) regulates mRNA stability of *BACE1*, a key enzyme in Alzheimer's disease pathology [10]. This subsequently affects amyloid- β 1-42 abundance, the increased expression of which is a hallmark of Alzheimer's disease. One mechanism by which lncRNAs have been hypothesized to impact neurodegenerative disease is through their induction of R-loop formation (which may be triggered by trinucleotide repeat expansion). R-loops have been shown to be capable of controlling the fate of neuroprotective genes [69], and are thought to contribute to the pathogenesis of fragile X syndrome and Friedrich's Ataxia [70, 71] by their silencing of certain genes. Additionally, work in *S. pombe* and *Arabidopsis* has suggested that R-loops may regulate lncRNA expression [72, 73], although whether this is true of lncRNAs linked to neurodegenerative diseases remains unclear. Trinucleotide repeats in lncRNAs are also known to be important in the pathogenesis of SCA8, by production of toxic noncoding CUG expansion RNAs from the ataxin 8 opposite strand (*ATXN8OS*), thought to cause a toxic gain of function at both the RNA and protein level [74, 75].

Another area of disease in which lncRNAs have been proven relevant is cardiovascular disease [66, 67]. Evidence now shows that lncRNAs are an important factor in susceptibility to coronary artery disease and myocardial infarction, prognosis in recovery from myocardial infarction, cardiovascular disease mortality, and heart failure [67]. Once again their correlations with prognosis and susceptibility have placed lncRNAs in the spotlight as a promising avenue of investigation in finding novel biomarkers.

Interestingly, *Drosophila* lncRNAs have been shown hold functional roles very relevant to these pathologies. *Hsromega* [76-80] and *bft* [81] are required for proper apoptosis process and cell differentiation, *yar* [82] and *CRG* [83] serve regulatory roles in the nervous system, and *scfA* and *scfB* are required for normal calcium transients and cardiac muscle contractility [19]. This is particularly promising given that these links can be made from the limited pool of *Drosophila* lncRNAs that have been experimentally characterised.

Molecular functions of lncRNA conserved in *Drosophila*

lncRNAs have been shown to function via a wide range of molecular mechanisms, falling under the broad categories of signals, molecular decoys, guide RNAs, or scaffolds [84]. Some lncRNAs have convincingly been shown to be translated, with the small peptide products (smORFs) having important biological functions [14-19, 22-25]. Through these various mechanisms (Figure 1), they have been implicated in

187 regulation of a diverse array of processes, such as differentiation, development, cell proliferation, nervous
188 system function, and cardiovascular function in both *Drosophila* and humans, despite the lack of sequence
189 conservation in lncRNAs across species. Importantly, similarities in the modes of action of lncRNAs have
190 been found at the molecular level between organisms, discussed below.

191 192 **lncRNA in the regulation of chromatin structure and gene expression**

193
194 One of the most extensively studied molecular mechanisms of lncRNA modes of action is their role in sex
195 chromosome dosage compensation pathways. Due to the difference in the number of X chromosome
196 copies between males and females, there exists a compensation pathway required to maintain a similar
197 level of expression for genes located on the X chromosome. In *Drosophila*, this is achieved by
198 transcriptional hyperactivation of the single copy of the genes in males, allowing their expression at
199 comparable levels to that given by the two copies of the gene found in females [85]. In humans, by
200 contrast, the genes located on the X-chromosome in human females are partially transcriptionally
201 repressed, giving a similar level of expression to that seen in males [86].

202
203 In *Drosophila*, the *RNA on the X* genes, *roX1* and *roX2*, are expressed in males, and regulate the assembly of
204 the Male Specific Lethal (MSL) complex in *Drosophila*; a chromatin modifier that functions in histone
205 modification [87-90]. The recruitment and binding of MSL proteins by high affinity sequences on the
206 nascent *roX* transcripts covering the X chromosome allows the assembly of the active MSL complex, which
207 can then spread in cis, allowing chromatin restructuring and hyperactivation of specific regions of the
208 chromosome.

209
210 An immediate comparison can be made between the *roX* genes in *Drosophila*, and lncRNAs involved in the
211 sex chromosome dosage compensation pathway in humans and other mammals; *X-inactive specific*
212 *transcript* (*Xist*) and its antisense transcript, *Tsix*. Like the *roX* genes, *Xist* coats the X chromosome, where it
213 regulates chromatin modifications, with consequent effects on the expression of particular target genes
214 [91, 92]. Unlike *roX*, *Xist* is expressed in females, and regulates the inactivation of the X chromosome by
215 facilitating the initiation and stabilising of the X chromosome inactivation process [86].

216
217 Although these lncRNA genes differ in their sequence, there are striking similarities between their role in
218 specific regulation of the X-chromosome and the molecular mechanisms by which they are thought to
219 achieve this. Interestingly, a subset of lncRNAs involved in chromatin looping, called topological anchor
220 point RNAs (tapRNAs), have been identified in the human and mouse genomes, with conserved zinc-finger
221 motifs capable of binding DNA and RNA [93]. Whether these are conserved in *Drosophila* has not yet been
222 studied, but given the involvement of lncRNAs in *Drosophila* chromatin regulation so far, this may be a
223 promising avenue to explore, and may reveal a wider conservation of this class of lncRNA chromatin
224 regulators.

225 226 **lncRNAs in the production of small peptides**

227
228 The *Drosophila sarcolamban (scl)* gene, originally classified as a lncRNA *pncr003* [94], is transcribed into a
229 992 base-pair mRNA, which is translated to produce two related peptides of less than 30 amino acids [19].
230 The *scl* gene is expressed in muscle cells, and *scl* null mutants show arrhythmic cardiac contractions, a
231 phenotype produced by abnormal intracellular calcium levels in contracting muscle cells [19].

232

Interestingly, the *scl* genes were found to have homologues in humans, namely *sarcolipin* (*sln*) and its longer paralogue, *phospholamban* (*pln*), encoding peptides of 31 and 52 amino acids respectively [19]. Phylogenetic analysis suggests that these genes belong to the same gene family, derived from a single ancestral gene, conserved for more than 550 million years. Furthermore, their function also seems to be conserved, with *Sln* and *Pln* regulating calcium transport in mammalian muscle cells, via dampening of Sarco-endoplasmic Reticulum Ca^{2+} adenosine triphosphate (SERCA) pump function. *Scl* peptides were able to colocalise and interact with *Drosophila* SERCA. Exogenous expression of the human *Pln* and *Sln* peptides in *Drosophila scl* mutant muscle cells were sufficient to rescue muscle function. Importantly, aberrant levels of *Sln* in humans have been linked to heart arrhythmias [95]. Regulation of SERCA by micropeptides (encoded by lncRNAs) has been extensively exploited in mammals; with tissue specific positive and negative regulators being found [22, 96, 97]. In addition, the number of characterized lncRNA genes encoding micropeptides is rapidly increasing, with roles found in a myriad of essential, conserved cellular functions, from phagocytosis [17] and cellular motility [98] to RNA degradation [18]. Thus, these examples show that lncRNAs that produce biologically relevant peptides may be conserved in structure, function, and relevance to pathologies between humans and *Drosophila* [19, 22].

Future directions

As previously shown in *sarcolamban*, proving the protein-coding potential of lncRNAs is a painstaking process, and an extremely difficult topic to broach; with genes having previously been catalogued as “non-coding” by arbitrary rules. Definitively showing the translation, or lack thereof, of an RNA using experimental techniques can be an arduous process, making this approach impractical to apply to the entire catalogue of identified lncRNAs. Ribosome profiling (in which a protease digestion is used to degrade RNA not protected by a bound ribosome,) and polysome profiling (where RNAs are separated by the number of ribosomes that are attached to different transcripts) have been used to provide a translational snapshot for several lncRNAs so far. This data has given a profile for lncRNA translation, but the threshold for significant translation is difficult to define in a non-arbitrary fashion. Therefore, use of model organisms to determine the biological function of any particular lncRNA remains crucial to gaining a meaningful understanding of the function of these molecules. A thorough and processive approach to clarifying this aspect of the gene class, as well as standardising measures and cut-offs for translational activity is an important priority for those in the field.

Bioinformatic approaches to elucidating the possible biological functions of lncRNAs are also being developed, although this method is not without its difficulties. Due to the poor sequence conservation characteristic of lncRNAs, standard approaches used to identify biologically relevant transcripts by their conservation within and across species are significantly less effective within this gene class. However, recent work has noted distinctive selection patterns in lncRNAs based on secondary structure [99], which may be of help in future analyses.

To conclude, we suggest that the studies currently being carried out on lncRNA in *Drosophila* should be of interest to a far wider audience than just fly geneticists, having shown that as a model organism, *Drosophila* is a logical choice both for better characterising this gene class, and for precursor studies to highlight genes and mechanisms that can be carried forward into more expensive and laborious large animal and human work. The superb annotation of the *Drosophila* genome and transcriptome, coupled with further increases in RNA-sequencing data available, will provide a candidate pool of lncRNAs for a rapid functional characterization (using the sophisticated genetic tools available in *Drosophila*). Therefore, further lncRNA

279 studies in *Drosophila*, of a suitably high calibre, are likely to provide us not only with a better understanding
280 of the basic science behind this gene class, but promise to highlight potential biomarkers, elucidate genetic
281 mechanisms behind a range of diseases, and perhaps provide novel targets for next generation
282 therapeutics.

283

284 **Abbreviations**

285

286 lncRNA, long non-coding RNA; MSL, Male Specific Lethal; pln, phospholamban; PRC2, Polycomb Repressive
287 Complex 2; roX, RNA on the X; scl, sarcolamban; sln, sarcolipin; small open reading frame, smORF; tapRNA,
288 topological anchor point RNA; Xist, X-inactive specific transcript.

289

290 **Funding**

291

292 This work was funded by a Brighton and Sussex Medical School studentship [grant number WB002-61] and
293 the Biotechnology and Biological Sciences Research Council [grant number BB/N001753/1].

294

295 **Competing Interests**

296

297 The Authors declare that there are no competing interests associated with the manuscript.

298

299 **Acknowledgements**

300

301 The authors thank Benjamin Towler and Pedro Patraquim for their input and advice.

302

References

- 1 Brown, C. J., Ballabio, A., Rupert, J. L., Lafreniere, R. G., Grompe, M., Tonlorenzi, R. and Willard, H. F. (1991) A gene from the region of the human X inactivation centre is expressed exclusively from the inactive X chromosome. *Nature*. **349**, 38-44
- 2 Taft, R. J., Pheasant, M. and Mattick, J. S. (2007) The relationship between non-protein-coding DNA and eukaryotic complexity. *Bioessays*. **29**, 288-299
- 3 Derrien, T., Johnson, R., Bussotti, G., Tanzer, A., Djebali, S., Tilgner, H., Guernec, G., Martin, D., Merkel, A., Knowles, D. G., Lagarde, J., Veeravalli, L., Ruan, X., Ruan, Y., Lassmann, T., Carninci, P., Brown, J. B., Lipovich, L., Gonzalez, J. M., Thomas, M., Davis, C. A., Shiekhatter, R., Gingeras, T. R., Hubbard, T. J., Notredame, C., Harrow, J. and Guigó, R. (2012) The GENCODE v7 catalog of human long noncoding RNAs: analysis of their gene structure, evolution, and expression. *Genome Res*. **22**, 1775-1789
- 4 Au, P. C., Zhu, Q. H., Dennis, E. S. and Wang, M. B. (2011) Long non-coding RNA-mediated mechanisms independent of the RNAi pathway in animals and plants. *RNA Biol*. **8**, 404-414
- 5 Hirota, K., Miyoshi, T., Kugou, K., Hoffman, C. S., Shibata, T. and Ohta, K. (2008) Stepwise chromatin remodelling by a cascade of transcription initiation of non-coding RNAs. *Nature*. **456**, 130-134
- 6 Tian, D., Sun, S. and Lee, J. T. (2010) The long noncoding RNA, Jpx, is a molecular switch for X chromosome inactivation. *Cell*. **143**, 390-403
- 7 Yoo, E. J., Cooke, N. E. and Liebhaber, S. A. (2012) An RNA-independent linkage of noncoding transcription to long-range enhancer function. *Mol Cell Biol*. **32**, 2020-2029
- 8 Lai, F., Orom, U. A., Cesaroni, M., Beringer, M., Taatjes, D. J., Blobel, G. A. and Shiekhatter, R. (2013) Activating RNAs associate with Mediator to enhance chromatin architecture and transcription. *Nature*. **494**, 497-501
- 9 Yoon, J. H., Abdelmohsen, K., Srikantan, S., Yang, X., Martindale, J. L., De, S., Huarte, M., Zhan, M., Becker, K. G. and Gorospe, M. (2012) LincRNA-p21 suppresses target mRNA translation. *Mol Cell*. **47**, 648-655
- 10 Faghihi, M. A., Modarresi, F., Khalil, A. M., Wood, D. E., Sahagan, B. G., Morgan, T. E., Finch, C. E., St Laurent, G., Kenny, P. J. and Wahlestedt, C. (2008) Expression of a noncoding RNA is elevated in Alzheimer's disease and drives rapid feed-forward regulation of beta-secretase. *Nat Med*. **14**, 723-730
- 11 Wang, H., Iacoangeli, A., Lin, D., Williams, K., Denman, R. B., Hellen, C. U. and Tiedge, H. (2005) Dendritic BC1 RNA in translational control mechanisms. *J Cell Biol*. **171**, 811-821
- 12 Gong, C. and Maquat, L. E. (2011) lncRNAs transactivate STAU1-mediated mRNA decay by duplexing with 3' UTRs via Alu elements. *Nature*. **470**, 284-288
- 13 Tripathi, V., Ellis, J. D., Shen, Z., Song, D. Y., Pan, Q., Watt, A. T., Freier, S. M., Bennett, C. F., Sharma, A., Bubulya, P. A., Blencowe, B. J., Prasanth, S. G. and Prasanth, K. V. (2010) The nuclear-retained noncoding RNA MALAT1 regulates alternative splicing by modulating SR splicing factor phosphorylation. *Mol Cell*. **39**, 925-938
- 14 Galindo, M. I., Pueyo, J. I., Fouix, S., Bishop, S. A. and Couso, J. P. (2007) Peptides encoded by short ORFs control development and define a new eukaryotic gene family. *PLoS Biol*. **5**, e106
- 15 Kondo, T., Hashimoto, Y., Kato, K., Inagaki, S., Hayashi, S. and Kageyama, Y. (2007) Small peptide regulators of actin-based cell morphogenesis encoded by a polycistronic mRNA. *Nat Cell Biol*. **9**, 660-665
- 16 Pueyo, J. I. and Couso, J. P. (2008) The 11-aminoacid long Tarsal-less peptides trigger a cell signal in *Drosophila* leg development. *Dev Biol*. **324**, 192-201
- 17 Pueyo, J. I., Magny, E. G., Sampson, C. J., Amin, U., Evans, I. R., Bishop, S. A. and Couso, J. P. (2016) Hemotin, a Regulator of Phagocytosis Encoded by a Small ORF and Conserved across Metazoans. *PLoS Biol*. **14**, e1002395
- 18 D'Lima, N. G., Ma, J., Winkler, L., Chu, Q., Loh, K. H., Corpuz, E. O., Budnik, B. A., Lykke-Andersen, J., Saghatelian, A. and Slavoff, S. A. (2017) A human microprotein that interacts with the mRNA decapping complex. *Nat Chem Biol*. **13**, 174-180
- 19 Magny, E. G., Pueyo, J. I., Pearl, F. M., Cespedes, M. A., Niven, J. E., Bishop, S. A. and Couso, J. P. (2013) Conserved regulation of cardiac calcium uptake by peptides encoded in small open reading frames. *Science*. **341**, 1116-1120

- 20 Laressergues, D., Couzigou, J. M., Clemente, H. S., Martinez, Y., Dunand, C., Bécard, G. and Combier, J. P. (2015) Primary transcripts of microRNAs encode regulatory peptides. *Nature*. **520**, 90-93
- 21 Williamson, L., Saponaro, M., Boeing, S., East, P., Mitter, R., Kantidakis, T., Kelly, G. P., Lobley, A., Walker, J., Spencer-Dene, B., Howell, M., Stewart, A. and Svejstrup, J. Q. (2017) UV Irradiation Induces a Non-coding RNA that Functionally Opposes the Protein Encoded by the Same Gene. *Cell*. **168**, 843-855.e813
- 22 Anderson, D. M., Anderson, K. M., Chang, C. L., Makarewich, C. A., Nelson, B. R., McAnally, J. R., Kasaragod, P., Shelton, J. M., Liou, J., Bassel-Duby, R. and Olson, E. N. (2015) A micropeptide encoded by a putative long noncoding RNA regulates muscle performance. *Cell*. **160**, 595-606
- 23 Aspden, J. L., Eyre-Walker, Y. C., Phillips, R. J., Amin, U., Mumtaz, M. A., Brocard, M. and Couso, J. P. (2014) Extensive translation of small Open Reading Frames revealed by Poly-Ribo-Seq. *Elife*. **3**, e03528
- 24 Mackowiak, S. D., Zuber, H., Bielow, C., Thiel, D., Kutz, K., Calviello, L., Mastrobuoni, G., Rajewsky, N., Kempa, S., Selbach, M. and Obermayer, B. (2015) Extensive identification and analysis of conserved small ORFs in animals. *Genome Biol*. **16**, 179
- 25 Ruiz-Orera, J., Messegue, X., Subirana, J. A. and Alba, M. M. (2014) Long non-coding RNAs as a source of new peptides. *Elife*. **3**, e03523
- 26 Reiter, L. T., Potocki, L., Chien, S., Gribskov, M. and Bier, E. (2001) A systematic analysis of human disease-associated gene sequences in *Drosophila melanogaster*. *Genome Res*. **11**, 1114-1125
- 27 Bier, E. (2005) *Drosophila*, the golden bug, emerges as a tool for human genetics. *Nat Rev Genet*. **6**, 9-23
- 28 Schoenfelder, S., Smits, G., Fraser, P., Reik, W. and Paro, R. (2007) Non-coding transcripts in the H19 imprinting control region mediate gene silencing in transgenic *Drosophila*. *EMBO Rep*. **8**, 1068-1073
- 29 Mercer, T. R. and Mattick, J. S. (2013) Structure and function of long noncoding RNAs in epigenetic regulation. *Nat Struct Mol Biol*. **20**, 300-307
- 30 Haerty, W. and Ponting, C. P. (2013) Mutations within lncRNAs are effectively selected against in fruitfly but not in human. *Genome Biol*. **14**, R49
- 31 Quinn, J. J., Zhang, Q. C., Georgiev, P., Ilik, I. A., Akhtar, A. and Chang, H. Y. (2016) Rapid evolutionary turnover underlies conserved lncRNA-genome interactions. *Genes Dev*. **30**, 191-207
- 32 Nguyen, D., Krueger, B. J., Sedore, S. C., Brogie, J. E., Rogers, J. T., Rajendra, T. K., Saunders, A., Matera, A. G., Lis, J. T., Uguen, P. and Price, D. H. (2012) The *Drosophila* 7SK snRNP and the essential role of dHEXIM in development. *Nucleic Acids Res*. **40**, 5283-5297
- 33 Ulitsky, I. (2016) Evolution to the rescue: using comparative genomics to understand long non-coding RNAs. *Nat Rev Genet*. **17**, 601-614
- 34 Cabili, M. N., Dunagin, M. C., McClanahan, P. D., Biaesch, A., Padovan-Merhar, O., Regev, A., Rinn, J. L. and Raj, A. (2015) Localization and abundance analysis of human lncRNAs at single-cell and single-molecule resolution. *Genome Biol*. **16**, 20
- 35 Lennox, K. A. and Behlke, M. A. (2016) Cellular localization of long non-coding RNAs affects silencing by RNAi more than by antisense oligonucleotides. *Nucleic Acids Res*. **44**, 863-877
- 36 Ji, P., Diederichs, S., Wang, W., Böing, S., Metzger, R., Schneider, P. M., Tidow, N., Brandt, B., Buerger, H., Bulk, E., Thomas, M., Berdel, W. E., Serve, H. and Müller-Tidow, C. (2003) MALAT-1, a novel noncoding RNA, and thymosin beta4 predict metastasis and survival in early-stage non-small cell lung cancer. *Oncogene*. **22**, 8031-8041
- 37 Schmidt, L. H., Spieker, T., Koschmieder, S., Schäffers, S., Humberg, J., Jungen, D., Bulk, E., Hascher, A., Wittmer, D., Marra, A., Hillejan, L., Wiebe, K., Berdel, W. E., Wiewrodt, R. and Muller-Tidow, C. (2011) The long noncoding MALAT-1 RNA indicates a poor prognosis in non-small cell lung cancer and induces migration and tumor growth. *J Thorac Oncol*. **6**, 1984-1992
- 38 Tano, K., Mizuno, R., Okada, T., Rakwal, R., Shibato, J., Masuo, Y., Ijiri, K. and Akimitsu, N. (2010) MALAT-1 enhances cell motility of lung adenocarcinoma cells by influencing the expression of motility-related genes. *FEBS Lett*. **584**, 4575-4580
- 39 Ren, S., Liu, Y., Xu, W., Sun, Y., Lu, J., Wang, F., Wei, M., Shen, J., Hou, J., Gao, X., Xu, C., Huang, J. and Zhao, Y. (2013) Long noncoding RNA MALAT-1 is a new potential therapeutic target for castration resistant prostate cancer. *J Urol*. **190**, 2278-2287
- 40 Hao, Y., Crenshaw, T., Moulton, T., Newcomb, E. and Tycko, B. (1993) Tumour-suppressor activity of H19 RNA. *Nature*. **365**, 764-767

408 41 Li, H., Yu, B., Li, J., Su, L., Yan, M., Zhu, Z. and Liu, B. (2014) Overexpression of lncRNA H19 enhances
409 carcinogenesis and metastasis of gastric cancer. *Oncotarget*. **5**, 2318-2329

410 42 Yang, F., Bi, J., Xue, X., Zheng, L., Zhi, K., Hua, J. and Fang, G. (2012) Up-regulated long non-coding
411 RNA H19 contributes to proliferation of gastric cancer cells. *FEBS J*. **279**, 3159-3165

412 43 Kogo, R., Shimamura, T., Mimori, K., Kawahara, K., Imoto, S., Sudo, T., Tanaka, F., Shibata, K.,
413 Suzuki, A., Komune, S., Miyano, S. and Mori, M. (2011) Long noncoding RNA HOTAIR regulates polycomb-
414 dependent chromatin modification and is associated with poor prognosis in colorectal cancers. *Cancer Res*.
415 **71**, 6320-6326

416 44 Wu, Z. H., Wang, X. L., Tang, H. M., Jiang, T., Chen, J., Lu, S., Qiu, G. Q., Peng, Z. H. and Yan, D. W.
417 (2014) Long non-coding RNA HOTAIR is a powerful predictor of metastasis and poor prognosis and is
418 associated with epithelial-mesenchymal transition in colon cancer. *Oncol Rep*. **32**, 395-402

419 45 Pang, Q., Ge, J., Shao, Y., Sun, W., Song, H., Xia, T., Xiao, B. and Guo, J. (2014) Increased expression
420 of long intergenic non-coding RNA LINC00152 in gastric cancer and its clinical significance. *Tumour Biol*. **35**,
421 5441-5447

422 46 Dong, Y., Liang, G., Yuan, B., Yang, C., Gao, R. and Zhou, X. (2015) MALAT1 promotes the
423 proliferation and metastasis of osteosarcoma cells by activating the PI3K/Akt pathway. *Tumour Biol*. **36**,
424 1477-1486

425 47 Cai, X., Liu, Y., Yang, W., Xia, Y., Yang, C., Yang, S. and Liu, X. (2016) Long noncoding RNA MALAT1 as
426 a potential therapeutic target in osteosarcoma. *J Orthop Res*. **34**, 932-941

427 48 Wang, J. Z., Xu, C. L., Wu, H. and Shen, S. J. (2017) LncRNA SNHG12 promotes cell growth and
428 inhibits cell apoptosis in colorectal cancer cells. *Braz J Med Biol Res*. **50**, e6079

429 49 Sun, J., Wang, X., Fu, C., Zou, J., Hua, H. and Bi, Z. (2016) Long noncoding RNA FGFR3-AS1 promotes
430 osteosarcoma growth through regulating its natural antisense transcript FGFR3. *Mol Biol Rep*. **43**, 427-436

431 50 Cong, M., Li, J., Jing, R. and Li, Z. (2016) Long non-coding RNA tumor suppressor candidate 7
432 functions as a tumor suppressor and inhibits proliferation in osteosarcoma. *Tumour Biol*. **37**, 9441-9450

433 51 Ma, B., Li, M., Zhang, L., Huang, M., Lei, J. B., Fu, G. H., Liu, C. X., Lai, Q. W., Chen, Q. Q. and Wang,
434 Y. L. (2016) Upregulation of long non-coding RNA TUG1 correlates with poor prognosis and disease status in
435 osteosarcoma. *Tumour Biol*. **37**, 4445-4455

436 52 Li, F., Cao, L., Hang, D., Wang, F. and Wang, Q. (2015) Long non-coding RNA HOTTIP is up-regulated
437 and associated with poor prognosis in patients with osteosarcoma. *Int J Clin Exp Pathol*. **8**, 11414-11420

438 53 Marques Howarth, M., Simpson, D., Ngok, S. P., Nieves, B., Chen, R., Siprashvili, Z., Vaka, D., Breese,
439 M. R., Crompton, B. D., Alexe, G., Hawkins, D. S., Jacobson, D., Brunner, A. L., West, R., Mora, J., Stegmaier,
440 K., Khavari, P. and Sweet-Cordero, E. A. (2014) Long noncoding RNA EWSAT1-mediated gene repression
441 facilitates Ewing sarcoma oncogenesis. *J Clin Invest*. **124**, 5275-5290

442 54 Wang, Y., Yao, J., Meng, H., Yu, Z., Wang, Z., Yuan, X., Chen, H. and Wang, A. (2015) A novel long
443 non-coding RNA, hypoxia-inducible factor-2 α promoter upstream transcript, functions as an inhibitor of
444 osteosarcoma stem cells in vitro. *Mol Med Rep*. **11**, 2534-2540

445 55 Min, L., Hong, S., Duan, H., Zhou, Y., Zhang, W., Luo, Y., Shi, R. and Tu, C. (2016) Antidifferentiation
446 Noncoding RNA Regulates the Proliferation of Osteosarcoma Cells. *Cancer Biother Radiopharm*. **31**, 52-57

447 56 Ruan, W., Wang, P., Feng, S., Xue, Y. and Li, Y. (2016) Long non-coding RNA small nucleolar RNA
448 host gene 12 (SNHG12) promotes cell proliferation and migration by upregulating angiomin gene
449 expression in human osteosarcoma cells. *Tumour Biol*. **37**, 4065-4073

450 57 Esteller, M. (2011) Non-coding RNAs in human disease. *Nat Rev Genet*. **12**, 861-874

451 58 Li, L., Chen, H., Gao, Y., Wang, Y. W., Zhang, G. Q., Pan, S. H., Ji, L., Kong, R., Wang, G., Jia, Y. H., Bai,
452 X. W. and Sun, B. (2016) Long Noncoding RNA MALAT1 Promotes Aggressive Pancreatic Cancer Proliferation
453 and Metastasis via the Stimulation of Autophagy. *Mol Cancer Ther*. **15**, 2232-2243

454 59 Meller, V. H., Joshi, S. S. and Deshpande, N. (2015) Modulation of Chromatin by Noncoding RNA.
455 *Annu Rev Genet*. **49**, 673-695

456 60 Roberts, T. C., Morris, K. V. and Wood, M. J. (2014) The role of long non-coding RNAs in
457 neurodevelopment, brain function and neurological disease. *Philos Trans R Soc Lond B Biol Sci*. **369**

458 61 Mori, K., Arzberger, T., Grässer, F. A., Gijssels, I., May, S., Rentzsch, K., Weng, S. M., Schludi, M.
459 H., van der Zee, J., Cruts, M., Van Broeckhoven, C., Kremmer, E., Kretschmar, H. A., Haass, C. and Edbauer,

D. (2013) Bidirectional transcripts of the expanded C9orf72 hexanucleotide repeat are translated into aggregating dipeptide repeat proteins. *Acta Neuropathol.* **126**, 881-893
 62 Zu, T., Liu, Y., Bañez-Coronel, M., Reid, T., Pletnikova, O., Lewis, J., Miller, T. M., Harms, M. B.,
 63 Falchook, A. E., Subramony, S. H., Ostrow, L. W., Rothstein, J. D., Troncoso, J. C. and Ranum, L. P. (2013)
 64 RAN proteins and RNA foci from antisense transcripts in C9ORF72 ALS and frontotemporal dementia. *Proc*
 65 *Natl Acad Sci U S A.* **110**, E4968-4977
 66 Lee, D. Y., Moon, J., Lee, S. T., Jung, K. H., Park, D. K., Yoo, J. S., Sunwoo, J. S., Byun, J. I., Shin, J. W.,
 67 Jeon, D., Jung, K. Y., Kim, M., Lee, S. K. and Chu, K. (2015) Distinct Expression of Long Non-Coding RNAs in
 68 an Alzheimer's Disease Model. *J Alzheimers Dis.* **45**, 837-849
 69 Johnson, R., Teh, C. H., Jia, H., Vanisri, R. R., Pandey, T., Lu, Z. H., Buckley, N. J., Stanton, L. W. and
 70 Lipovich, L. (2009) Regulation of neural macroRNAs by the transcriptional repressor REST. *RNA.* **15**, 85-96
 71 Johnson, R. (2012) Long non-coding RNAs in Huntington's disease neurodegeneration. *Neurobiol*
 72 *Dis.* **46**, 245-254
 73 Uchida, S. and Dimmeler, S. (2015) Long noncoding RNAs in cardiovascular diseases. *Circ Res.* **116**,
 74 737-750
 75 Archer, K., Broskova, Z., Bayoumi, A. S., Teoh, J. P., Davila, A., Tang, Y., Su, H. and Kim, I. M. (2015)
 76 Long Non-Coding RNAs as Master Regulators in Cardiovascular Diseases. *Int J Mol Sci.* **16**, 23651-23667
 77 Portoso, M., Ragazzini, R., Brenčič, Ž., Moiani, A., Michaud, A., Vassilev, I., Wassef, M., Servant, N.,
 78 Sargueil, B. and Margueron, R. (2017) PRC2 is dispensable for HOTAIR-mediated transcriptional repression.
 79 *EMBO J.* **36**, 981-994
 80 Salvi, J. S. and Mekhail, K. (2015) R-loops highlight the nucleus in ALS. *Nucleus.* **6**, 23-29
 81 Groh, M., Lufino, M. M., Wade-Martins, R. and Gromak, N. (2014) R-loops associated with triplet
 82 repeat expansions promote gene silencing in Friedreich ataxia and fragile X syndrome. *PLoS Genet.* **10**,
 83 e1004318
 84 Colak, D., Zaninovic, N., Cohen, M. S., Rosenwaks, Z., Yang, W. Y., Gerhardt, J., Disney, M. D. and
 85 Jaffrey, S. R. (2014) Promoter-bound trinucleotide repeat mRNA drives epigenetic silencing in fragile X
 86 syndrome. *Science.* **343**, 1002-1005
 87 Sun, Q., Csorba, T., Skourti-Stathaki, K., Proudfoot, N. J. and Dean, C. (2013) R-loop stabilization
 88 represses antisense transcription at the Arabidopsis FLC locus. *Science.* **340**, 619-621
 89 Nakama, M., Kawakami, K., Kajitani, T., Urano, T. and Murakami, Y. (2012) DNA-RNA hybrid
 90 formation mediates RNAi-directed heterochromatin formation. *Genes Cells.* **17**, 218-233
 91 Tan, H., Xu, Z. and Jin, P. (2012) Role of noncoding RNAs in trinucleotide repeat neurodegenerative
 92 disorders. *Exp Neurol.* **235**, 469-475
 93 Mutsuddi, M., Marshall, C. M., Benzow, K. A., Koob, M. D. and Rebay, I. (2004) The spinocerebellar
 94 ataxia 8 noncoding RNA causes neurodegeneration and associates with staufen in Drosophila. *Curr Biol.* **14**,
 95 302-308
 96 Lakhota, S. C., Mallik, M., Singh, A. K. and Ray, M. (2012) The large noncoding hsrw-n transcripts
 97 are essential for thermotolerance and remobilization of hnRNPs, HP1 and RNA polymerase II during
 98 recovery from heat shock in Drosophila. *Chromosoma.* **121**, 49-70
 99 Prasanth, K. V., Rajendra, T. K., Lal, A. K. and Lakhota, S. C. (2000) Omega speckles - a novel class of
 100 nuclear speckles containing hnRNPs associated with noncoding hsr-omega RNA in Drosophila. *J Cell Sci.* **113**
 101 **Pt 19**, 3485-3497
 102 Perrimon, N., Lanjuin, A., Arnold, C. and Noll, E. (1996) Zygotic lethal mutations with maternal
 103 effect phenotypes in Drosophila melanogaster. II. Loci on the second and third chromosomes identified by
 104 P-element-induced mutations. *Genetics.* **144**, 1681-1692
 105 Mallik, M. and Lakhota, S. C. (2009) The developmentally active and stress-inducible noncoding
 106 hsromega gene is a novel regulator of apoptosis in Drosophila. *Genetics.* **183**, 831-852
 107 Johnson, T. K., Cockerell, F. E. and McKechnie, S. W. (2011) Transcripts from the Drosophila heat-
 108 shock gene hsr-omega influence rates of protein synthesis but hardly affect resistance to heat knockdown.
 109 *Mol Genet Genomics.* **285**, 313-323
 110 Hardiman, K. E., Brewster, R., Khan, S. M., Deo, M. and Bodmer, R. (2002) The bereft gene, a
 111 potential target of the neural selector gene cut, contributes to bristle morphogenesis. *Genetics.* **161**, 231-
 112 247

513 82 Soshnev, A. A., Ishimoto, H., McAllister, B. F., Li, X., Wehling, M. D., Kitamoto, T. and Geyer, P. K.
 514 (2011) A conserved long noncoding RNA affects sleep behavior in *Drosophila*. *Genetics*. **189**, 455-468
 515 83 Li, M., Wen, S., Guo, X., Bai, B., Gong, Z., Liu, X., Wang, Y., Zhou, Y., Chen, X., Liu, L. and Chen, R.
 516 (2012) The novel long non-coding RNA CRG regulates *Drosophila* locomotor behavior. *Nucleic Acids Res.* **40**,
 517 11714-11727
 518 84 Wang, K. C. and Chang, H. Y. (2011) Molecular mechanisms of long noncoding RNAs. *Mol Cell*. **43**,
 519 904-914
 520 85 Deng, X. and Meller, V. H. (2006) roX RNAs are required for increased expression of X-linked genes
 521 in *Drosophila melanogaster* males. *Genetics*. **174**, 1859-1866
 522 86 Pontier, D. B. and Gribnau, J. (2011) Xist regulation and function explored. *Hum Genet.* **130**, 223-
 523 236
 524 87 Park, Y., Kelley, R. L., Oh, H., Kuroda, M. I. and Meller, V. H. (2002) Extent of chromatin spreading
 525 determined by roX RNA recruitment of MSL proteins. *Science*. **298**, 1620-1623
 526 88 Kelley, R. L., Lee, O. K. and Shim, Y. K. (2008) Transcription rate of noncoding roX1 RNA controls
 527 local spreading of the *Drosophila* MSL chromatin remodeling complex. *Mech Dev.* **125**, 1009-1019
 528 89 Kelley, R. L. and Kuroda, M. I. (2003) The *Drosophila* roX1 RNA gene can overcome silent chromatin
 529 by recruiting the male-specific lethal dosage compensation complex. *Genetics*. **164**, 565-574
 530 90 Oh, H., Park, Y. and Kuroda, M. I. (2003) Local spreading of MSL complexes from roX genes on the
 531 *Drosophila* X chromosome. *Genes Dev.* **17**, 1334-1339
 532 91 Plath, K., Mlynarczyk-Evans, S., Nusinow, D. A. and Panning, B. (2002) Xist RNA and the mechanism
 533 of X chromosome inactivation. *Annu Rev Genet.* **36**, 233-278
 534 92 McHugh, C. A., Chen, C. K., Chow, A., Surka, C. F., Tran, C., McDonel, P., Pandya-Jones, A., Blanco,
 535 M., Burghard, C., Moradian, A., Sweredoski, M. J., Shishkin, A. A., Su, J., Lander, E. S., Hess, S., Plath, K. and
 536 Guttman, M. (2015) The Xist lncRNA interacts directly with SHARP to silence transcription through HDAC3.
 537 *Nature*. **521**, 232-236
 538 93 Amaral, P. P., Leonardi, T., Han, N., Vire, E., Gascoigne, D. K., Arias-Carrasco, R., Zhang, A., Pluchino,
 539 S. and Maracaja-Coutinho, V. (2016) Genomic positional conservation identifies topological anchor point
 540 (tap) RNAs linked to developmental loci.
 541 94 Tupy, J. L., Bailey, A. M., Dailey, G., Evans-Holm, M., Siebel, C. W., Misra, S., Celniker, S. E. and
 542 Rubin, G. M. (2005) Identification of putative noncoding polyadenylated transcripts in *Drosophila*
 543 *melanogaster*. *Proc Natl Acad Sci U S A.* **102**, 5495-5500
 544 95 Shanmugam, M., Molina, C. E., Gao, S., Severac-Bastide, R., Fischmeister, R. and Babu, G. J. (2011)
 545 Decreased sarcolipin protein expression and enhanced sarco(endo)plasmic reticulum Ca²⁺ uptake in
 546 human atrial fibrillation. *Biochem Biophys Res Commun.* **410**, 97-101
 547 96 Anderson, D. M., Makarewich, C. A., Anderson, K. M., Shelton, J. M., Bezprozvannaya, S., Bassel-
 548 Duby, R. and Olson, E. N. (2016) Widespread control of calcium signaling by a family of SERCA-inhibiting
 549 micropeptides. *Sci Signal.* **9**, ra119
 550 97 Nelson, B. R., Makarewich, C. A., Anderson, D. M., Winders, B. R., Troupes, C. D., Wu, F., Reese, A.
 551 L., McAnally, J. R., Chen, X., Kavalali, E. T., Cannon, S. C., Houser, S. R., Bassel-Duby, R. and Olson, E. N.
 552 (2016) A peptide encoded by a transcript annotated as long noncoding RNA enhances SERCA activity in
 553 muscle. *Science*. **351**, 271-275
 554 98 Pauli, A., Norris, M. L., Valen, E., Chew, G. L., Gagnon, J. A., Zimmerman, S., Mitchell, A., Ma, J.,
 555 Dubrulle, J., Reyon, D., Tsai, S. Q., Joung, J. K., Saghatelian, A. and Schier, A. F. (2014) Toddler: an embryonic
 556 signal that promotes cell movement via Apelin receptors. *Science*. **343**, 1248636
 557 99 Pegueroles, C. and Gabaldón, T. (2016) Secondary structure impacts patterns of selection in human
 558 lncRNAs. *BMC Biol.* **14**, 60
 559
 560
 561

562 **Figures**

563
564 **Table 1) A table summarising the lncRNAs linked to various kinds of cancer, as covered in this review.**

lncRNA	Associated disease	Reference
MALAT1	Osteosarcoma	[46, 47]
	Non-small cell lung cancer	[36-38]
	Prostate cancer	[39]
	Pancreatic cancer	[58]
HOTAIR	Colorectal cancer	[43, 44]
EWSAT1	Ewing sarcoma	[53]
HOTTIP	Osteosarcoma	[52]
HIF2PUT	Osteosarcoma	[54]
ANCR	Osteosarcoma	[55]
TUSC7	Osteosarcoma	[50]
FGFR3-AS1	Osteosarcoma	[49]
SNHG12	Osteosarcoma	[56]
TUG1	Osteosarcoma	[51]
H19	Wilms tumour	[40]
	Gastric cancer	[41, 42]
LINC00152	Gastric cancer	[45]

567 **Figure 1) A cartoon depicting the molecular mechanisms by which lncRNAs can function.**

568 a) Some lncRNAs (red), such as *Xist* and *RoX1*, can act to modulate expression of a certain gene by binding
569 to a transcription modifier or chromatin modifier (purple). b) lncRNAs (red) such as *HOTAIR* can act as
570 molecular scaffolds, allowing the assembly of protein complexes (teal, green, dark purple) with genetic
571 regulatory roles e.g. polycomb complex PRC2. c) lncRNAs (red) can act as molecular decoys, to sequester
572 miRNAs (orange) or proteins (purple). d) Alternatively, lncRNAs (red) can act as molecular decoys, occluding
573 or removing transcription factors, proteins, or RNAs (purple) from their functional location. e) lncRNAs
574 (red) can act as a molecular guide, allowing formation of ribonucleoprotein complexes (yellow) to specific
575 target sites. f) It has also been shown that lncRNAs (blue as DNA, red as RNA) can be actively translated into
576 functional smORF peptides (orange) such as the *SclA* and *SclB* peptides, which function in regulating
577 calcium transport in cardiac muscle.

